

# Stationarity and Spatiotemporal Dynamics at the Orbital Ordering Transition in the Half-doped Manganite $\text{Pr}_{0.5}\text{Ca}_{0.5}\text{MnO}_3$

**Joshua Turner**<sup>1,2</sup>, Jessica Thomas<sup>3</sup>, John Hill<sup>3</sup>, Mark Pfeifer<sup>1,2</sup>, Karine Chesnel<sup>2\*</sup>, Y. Tokura<sup>4,5,6</sup>, Y. Tomioka<sup>4</sup>, S.D. Kevan<sup>1</sup>.

<sup>1</sup>Physics Dept., University of Oregon; <sup>2</sup>LBNL, Berkeley, CA; <sup>3</sup>CM Physics, Brookhaven National Lab; <sup>4</sup>Correlated Electron Research Center, National Institute of Advanced Industrial Science and Technology, Tsukuba, Japan; <sup>5</sup>Department of Applied Physics, University of Tokyo, Japan; <sup>6</sup>Spin Superstructure Project, ERATO, Japan Science and Technology Corporation, Tsukuma, Japan; \*now at LNCMP, Toulouse, France.

Understanding the mesoscale structure and dynamics of complex charge, spin, and orbital-ordered phases is crucial to unraveling the mystery of the colossal magnetoresistance (CMR) phenomenon. The use of coherent soft x-rays allows this complexity to be mapped into far-field speckle-diffraction patterns, which can be analyzed to probe spatiotemporal dynamics, microscopic memory effects, and real-space structures through phase retrieval. We report new coherent soft x-ray scattering experiments performed on beamline 12.0.2.2 at the Advanced Light Source wherein we probe the structure and dynamics of orbital domains in  $\text{Pr}_{0.5}\text{Ca}_{0.5}\text{MnO}_3$  near the orbital ordering transition temperature. By operating at the Bragg condition for the lowest-order orbital reflection and at a wavelength that is resonant with the Mn L-edge, we have obtained excellent scattering signal with direct sensitivity to the orbital ordered phase. We find that the low temperature state is characterized by static short-range ordered domains and that the system remains largely static as it is warmed through the disordering transition. In addition, we have observed small-amplitude spatiotemporal fluctuations near the transition. We have studied the connection between this small fluctuating component, a decrease in the orbital domain coherence length, and a change in shape of the integrated scattering profile over a narrow temperature range all just below the transition. Our measurements present new insight into how the orbital ordering domain state behaves near the order-disorder transformation - a clue into a key ingredient of the CMR story.